

---

## A Method to Quantify Biofilms in Object Glass Using ImageJ

Arief Heru<sup>1\*</sup>, Reza Hakim<sup>1</sup>, Hanggia Primadita<sup>2</sup>, and Rio Risandiansyah<sup>1</sup>

<sup>1</sup>Faculty of Medicine, University of Islam Malang, Malang 65144, Indonesia

<sup>2</sup>Faculty of Medicine, Brawijaya University, Malang 65145, Indonesia

\*Corresponding Author

Email : [a.heru2878@gmail.com](mailto:a.heru2878@gmail.com)

---

### Abstract

One method to observe biofilms is by inoculating bacteria on glass slides and observed for the presence of large structures (microcolonies) and small structures (clusters, aggregates, or single cells) under a microscope. Analysis of these structures using image processing software may provide a method to quantify biofilm production and degradation in glass slides. In this study, we use ImageJ to quantify the number and area percentage of microbial structures observable on a slide. This study is an experimental *in vitro* study. Biofilm production was done by submerging slides in petri dishes filled with Brain Heart Infusion with 2% sucrose (w/v) and inoculating it with bacteria. The petri dishes were incubated undisturbed for 48 hours at 37°C (n=3). Afterwards, the slides were removed and submerged in distilled water (Group 1) or detergent (Group 2) for 5 minutes before staining with 0.1% crystal violet and rewashed. The slides were then observed under a light microscope at 1000x and images from five fields of view were collected. ImageJ was then used to count the number of microcolonies (>15.000 µm<sup>2</sup>), aggregate cells (200 – 14.999 µm<sup>2</sup>), and single cells or cell clusters (1 – 199 µm<sup>2</sup>), and their area percentage. Welch's T-Test was performed using JASP version 0.18.3. Observation of slides shows microcolonies to be formed in Group 1, and no or little in Group 2. Based on ImageJ calculation, slides treated with distilled water had a biofilm consisting of an average 4.60 ± 2.41 microcolony number and an average percentage area of 39.97 ± 9.99%, 120.47 ± 32.31 (8.96 ± 3.19%) cell aggregates, and 415.06 ± 139.85 (1.39 ± 0.33%) single cells and cell clusters. Detergent application possibly showed biofilm breakdown, with a significant (p<0.001) reduction in microcolony percentage area to up to 99% (0.33 ± 0.68% remaining) and increased single cell number and percentage area to 1,754.93 ± 689.52 (5.27 ± 0.49%). ImageJ can be a valuable tool to quantify biofilm production in glass slides based on the number and percentage area of microcolonies, cell aggregates, and single cells or cell clusters.

**Keywords:** Biofilm, ImageJ, Direct Microscopic Observation, Quantification

---

## INTRODUCTION

Microbial biofilms are a current problem particularly in medical device-related infections, particularly from invasive medical devices such as catheters and endotracheal tubes (Dexter & Scott, 2019; Di Domenico et al., 2022). Biofilms are a constant source of bacterial nosocomial infection and major cause of septicemia or sepsis which increases the health burden in hospitals (Rubi et al., 2022). Furthermore, it is known that biofilms can also withstand antibiotic therapy, whether by the dissemination of antibacterial genes from persistent bacteria in a biofilm community, by reducing antibiotic exposure inside the biofilm due to the production of extracellular polysaccharide substances in the biofilm, or by the antibiotic unable to work on metabolically dormant cells located in the base of the biofilm (Yan & Bassler, 2019). Thus, efforts to screen for agents that can reduce biofilm, either by inhibiting its production or eradicating mature biofilm, are currently being undertaken worldwide. This effort requires methods to quantify and observe microbial biofilm *in vitro*.

There are different states of the microbial cells in a biofilm produced on an object glass, which also coincides with the current theory or observation of biofilm production. Biofilm production is initiated when a bacterium attaches to a surface and begins to replicate. This would form several cells as well as provide a platform for other cell recruitment during the initial phase of extracellular polymeric substance production (Schilcher & Horswill, 2020). After more cells are attached, a structural community of bacteria is established, which, as well as being comparatively large in size, also provides nutrition and means to communicate with other cells in this community (Di Domenico et al., 2022; Ruhai & Kataria, 2021). When viewed using a confocal laser scanning microscope (CLSM), it has been discovered that microbial biofilms on an apatitic surface would have various

configurations, consisting of microbial in single cells (around 1 – 5  $\mu\text{m}$ ), a cluster of cells (5 – 20  $\mu\text{m}$ ), an aggregate of cells (20 – 300  $\mu\text{m}$ ), or a large microcolony (>300  $\mu\text{m}$ )(Paula et al., 2020).

Several methods to produce and quantify biofilm in a laboratory setting has been explored in other papers(Coffey & Anderson, 2014); however, no golden standard for biofilm production and quantification has been established. In one method, an object glass is used as an abiotic surface for bacterial attachment and biofilm production and stained with crystal violet (usually 0.1%)(Bakkiyaraj et al., 2017; Katsipis et al., 2021). Crystal violet is reported to be able to stain cell walls as well as extracellular polymeric substances (EPS) which compose the biofilm matrix(Dertli et al., 2015), and therefore enable the visualization of biofilm-related microscopic structures.

Image files acquired from microscopic observation can be analyzed digitally by using image processing software. One of these, often used in various cell-based research projects, is ImageJ – a free java-based image processing software available for most operating systems. ImageJ can be used to calculate area, measure distances and angles, and create density histograms(Rueden et al., 2017). ImageJ has been used in many research studies to identify and calculate abnormal cell numbers, cell size and malformation(Young & Morrison, 2018), and bacterial cell counts on a plate(Schulze et al., 2011). However, the use of ImageJ in biofilm related research is limited.

This study aims to provide a novel method to quantify biofilms from images obtained using direct microscopic observation of biofilm produced in object glass. Firstly, we produced biofilm in object glass in two groups. In one group, the object glass was treated with detergent prior to staining, and the other was treated with water. We then use ImageJ to quantify biofilm based on the area size and the number of large structures (microcolonies and cell aggregates) and small structures (single cells) and compare the biofilm between those two groups.

## RESEARCH METHODS

### Bacteria Preparation

A clinical *Staphylococcus aureus* strain was used in this study, which was known to be able to produce biofilm. Bacterial was grown in a Nutrient Broth (Himedia) for 24 hours and plated in nutrient agar (Himedia) every month. For this study, several colonies were taken and mixed with normal saline and standardized using a 0.5 McFarland Standard, or equivalent to  $0.5 \times 10^8$  CFU/ml. However, contamination by an unknown basil bacteria occurred during the production of the biofilm, which occurred despite the production of biofilm on different slides on different days. The data collected by this unknown basil bacteria was used in this study.

### Biofilm Production and Staining Using Microscopic Direct Observation

Biofilm production was done according to the direct microscopy observation method as described in previous literature with several modifications(Bakkiyaraj et al., 2017), conducted with three biological replicates on different days. A sterilized glass object was inserted into the middle of a sterilized petri dish under aseptic conditions. Each petri dish was then filled with 10 ml of Brain Heart Infusion Broth (HiMedia) supplemented with 2% sucrose (w/v). From the standardized bacterial stocks, 100  $\mu\text{l}$  was transferred into the petri dish and was incubated undisturbed for 48 hours at 37°C. After incubation, washing was performed on each slide by submerging it in distilled water or detergent, respectively, for 5 minutes before staining. The object glass was transferred into a 0.1% Crystal Violet (CV) solution, left for 1 minute, and then washed by submerging in distilled water. The object glasses were then left to dry overnight before microscopic observation.

### Biofilm Observation and Quantification Using Imagej

Observation of the object glass was done on a trinocular microscope at 1000x magnification using emersion oil. From each object glass, five different fields of view were obtained and saved as a .TIFF image. Quantification was performed using ImageJ to determine the number of single cells, cell clusters, cell aggregates, and microcolonies, as well as the percentage area size.

ImageJ was used to analyze the images according to the manual(Rueden et al., 2017). Briefly,

each picture was transformed into an 8-bit image. The unit of measurement was changed into  $\mu\text{m}$ , based on the scale on each image (each image had a 200  $\mu\text{m}$  scale line) (using the 'set scale' function). The threshold was then adjusted to remove background, resulting in a black and white image. The function 'analyze particle' was used to calculate the number of structures detected, by inserting the area sizes for each structure, as described below. The circularity was left at default at 0.00 – 1.00. The option to 'include holes' was selected, and each result was summarized.

Prior to calculation, a sample microscopic image was used to calibrate the area size measurement used to identify microcolonies, aggregate cells, and single cells. The definition of each structure was obtained from previous literature; however, it should be noted that not all those structures were perfect circles. As far as we are aware, ImageJ calculates and identifies objects using area size, not diameter. Thus, the area determined was tweaked from preselected images so as to select only the intended structure, which fulfills the size requirement based on literature. The obtained area sizes for each structure were as follows: microcolonies ( $>15.000 \mu\text{m}^2$ ), aggregate cells (200 – 14.999  $\mu\text{m}^2$ ), and cluster cells or single cells (1 – 199  $\mu\text{m}^2$ ). The scale for each image was 1.13 pixels/ $\mu\text{m}$ .


### **Statistical Analysis**

Statistical analysis was conducted using JASP version 0.18.3 (JASP, 2024) using Welch's t-test, with a p-value considered to be  $< 0.05$ .

## **RESULTS AND DISCUSSION**

### **Microscopic Observation Showed Bacterial Microcolony and Cell Aggregate Formation on Slides**

Similar to previous literature (Bakkiyaraj et al., 2017; Katsipis et al., 2021), our results similarly show that microcolonies and cell clusters or aggregates were formed on the surface of the slides, as shown in Figure 1. Studies have stated that Crystal Violet (CV) can stain bacterial cell walls as well as extracellular polysaccharides (EPS) produced in biofilm formation (Giordani et al., 2023). However, any EPS staining was not observable in our findings. Bacterial cells were observed to be in close proximity to each other and forming a large structure. One limitation of this study, compared to previous studies in this subject, is the inability to observe three dimensional structures; thus, it is unknown whether the structure form is also constructed vertically as it would in a typical microcolony structure in biofilm production. However, as the diameter of the structure formed was well beyond 300  $\mu\text{m}$ , we conclude that the large structures observed were one or several microcolonies.



**Figure 1. Pre-processing (left) and post-processing (right) of images using Image-J to quantify biofilm production, with a blue overlay indicating a microcolony structure.**

Aside from microcolonies, smaller structures such as single cells and cell clusters were also visible. The definitions of a cell cluster and aggregate were a collection of cells with a diameter of 1–5  $\mu\text{m}$  and 5 – 50  $\mu\text{m}$ , respectively (Paula et al., 2020). However, using ImageJ, identification of structure was done using area size instead of diameter, and therefore, the selection based on structural diameter of these structures was difficult to obtain. Hence, in this paper, both cell clusters and single cells were grouped into one. This is done considering that aside from the difference in diameter, these two structures are not known to have a different role in the formation of biofilm – they can both be considered as transitional states between single cells, which are the singular or small cell structures that would initiate biofilm production, and the microcolony structure, which is often considered to be the main key structure in bacterial biofilms. On the other hand, it can be hypothesized that an increase in the number of these smaller structures, especially with the absence of large microcolonial structures, would indicate a breakdown of the current biofilm structure.


**Table 1. Summary of bacterial structures calculated using Image-J (three slides from different days with five fields of view per slide)**

No.	Parameters	Group 1 <sup>a</sup>	Group 2 <sup>a</sup>
1.	Average Single and Cluster cells		
	Number (cell units)	415.06 ± 139.85	1,754.93 ± 689.52*
	Percentage Area Size (%)	1.34 ± 0.33	5.27 ± 0.49*
2.	Average Aggregate		
	Number (cell units)	120.47 ± 32.31	193.34      73.87
	Percentage Area Size (%)	8.96 ± 3.19	7.22 ± 2.39
3.	Average Microcolony		
	Number (cell units)	4.60 ± 2.41	0.20      0.4*
	Percentage Area Size (%)	39.97 ± 9.99	0.33 ± 0.68*

<sup>a</sup>Group 1 was slides treated with water and Group 2 was slides treated with detergent. (\*) notation signifies a significant difference between Group 1 and Group 2.

### **Quantification of bacterial structures using ImageJ showed water-treated slides having a high percentage of microcolony and cell aggregate structures**


As shown in Figure 1, ImageJ was able to detect both microcolonies as well as smaller structures. The results of this calculation are shown in Table 1. In our case, slides washed with water (Group 1) showed <10 microcolonies per field of view at every slide, as the close proximity of the cells caused it to form one continuous structure. The microcolony had an average microcolony percentage area size of around  $39.97 \pm 9.99\%$  of the image, which represented the size of the microcolony. The number of clusters or aggregates, on the other hand, was around  $120.47 \pm 32.31$  structures per field of view for every slide, with a total area size of around  $4.60 \pm 2.41\%$ , and the single cells were  $415.06 \pm 139.85$  with a total area of  $1.34 \pm 0.33\%$ . Direct microscopic observation shows that detergent treatment was able to reduce the number of microcolonies observed in each slide, as shown in Figure 2. Quantification using ImageJ shows that slides treated with detergent showed a significant reduction in the microcolony number and area percentage, and an increase in single cell number and area percentage (Figure 3). However, no significant difference was observed in aggregate cell number or area percentage.



**Figure 2. A side-by-side comparison of typical microscopic images of different slides treated with water (left) and detergent (right). Little or no microcolonies were observed in most slides treated with detergent.**

Disinfectants are often used in laboratories for the sterilization of laboratory equipment. In the last 5 years, there has been an increasing interest in comparing the efficacy of disinfectants in removing biofilm (Osland et al., 2023). Based on our results, submerging in disinfectant was able to reduce the microcolony area by up to 99% - however, the number and percentage area of single cells increased after treatment. This may indicate that the detergent was able to breakdown microcolonies and cause

the destabilization of preformed biofilms. As the protocol used in this study attempts to reduce any mechanical cleansing of the slides (the slides are submerged and not passed through running water or detergent), we speculate that the cells in this protocol would remain on the surface of the slides instead of being displaced. In normal cleaning, this detergent would be effective in removing and sterilizing equipment from biofilms and bacterial cells.



**Figure 3. Bar-Plots comparing the number and area size percentage of single cells, cell aggregates, and microcolonies using Image-J on Group 1 (treated with water) and Group 2 (treated with detergent). Bars denote a confidence of interval 95%.**

Biofilm production is influenced by many different factors - different bacteria would have different biofilm production capabilities, the type of media used in biofilm production (as well as the addition of different sugars), and incubation time play a role in the success of biofilm production (Vandeplassche et al., 2017). This is usually crucial in the search for antibiofilm compounds – i.e., compounds that can prevent the production of biofilm by the bacteria. Some researchers screen for antibiofilm compounds against biofilm that are not yet established, whereas a compound of interest is added during the early stages of bacterial life (some times at the same time as the bacterial inoculation) (Mombeshora et al., 2021; Raissa et al., 2020). A more established biofilm would be harder to remove, as it would have defenses against chemical compounds through the formation of an extrapolymeric substance (EPS) layer that prevents any chemical compounds from interacting with the bacterial layer. Furthermore, bacteria located deeper in the biofilm structure would be metabolically inactive, preventing the metabolic action of certain antibacterial agents.

In this study, we show that Image J can be a valuable tool in determining biofilm production and size, as well as be used to measure antibiofilm bioactivity of certain compounds, and should be used in conjunction with direct microscopic observation for biofilm production. Increased biofilm production would be expected if a bacterial agent with high biofilm production was used as well as with optimized growth media and incubation time. As with the 96 well plate method, the number of washes used would influence final biofilm production, where increased washing would physically remove biofilm. Finally, as this method submerges the slides, a large volume of test compounds is required if this method is to be used as antibiofilm screening, therefore, this method might not be suitable for small sample volumes.

## CONCLUSIONS

ImageJ can be used to quantify biofilm production by calculating the number of and area size of microcolonies (area size of  $>15.000 \mu\text{m}^2$ ), cell aggregates and single cells. In biofilm, detergent

application was found to reduce the number of large structures and increase the number of smaller structures, showing a possible mechanism of biofilm breakdown. This method can be included in the repertoire of antibiofilm activity screening, and possibly be used to simulate biofilm production and destruction in clinically important mixed cultures found in various medical devices.

## REFERENCES

- Bakkiyaraj, D., Sritharadol, R., Padmavathi, A. R., Nakpheng, T., & Srichana, T. (2017). Antibiofilm properties of a mupirocin spray formulation against *Escherichia coli* wound infections. *Biofouling*, 33(7), 591–600. <https://doi.org/10.1080/08927014.2017.1337100>
- Coffey, B. M., & Anderson, G. G. (2014). Biofilm Formation in the 96-Well Microtiter Plate. In A. Filloux & J.-L. Ramos (Eds.), *Pseudomonas Methods and Protocols* (Vol. 1149, pp. 631–641). Springer New York. [https://doi.org/10.1007/978-1-4939-0473-0\\_48](https://doi.org/10.1007/978-1-4939-0473-0_48)
- Dertli, E., Mayer, M. J., & Narbad, A. (2015). Impact of the exopolysaccharide layer on biofilms, adhesion and resistance to stress in *Lactobacillus johnsonii* FI9785. *BMC Microbiology*, 15, 8. <https://doi.org/10.1186/s12866-015-0347-2>
- Dexter, A. M., & Scott, J. B. (2019). Airway management and ventilator-associated events. *Respiratory Care*, 64(8), 986–993. <https://doi.org/10.4187/respcare.07107>
- Di Domenico, E. G., Oliva, A., & Guembe, M. (2022). The current knowledge on the pathogenesis of tissue and medical device-related biofilm infections. *Microorganisms*, 10(7), 1259. <https://doi.org/10.3390/microorganisms10071259>
- Giordani, B., Naldi, M., Croatti, V., Parolin, C., Erdoğan, Ü., Bartolini, M., & Vitali, B. (2023). Exopolysaccharides from vaginal lactobacilli modulate microbial biofilms. *Microbial Cell Factories*, 22(1), 45. <https://doi.org/10.1186/s12934-023-02053-x>
- Katsipis, G., Tsaloukidou, V., Halevas, E., Geromichalou, E., Geromichalos, G., & Pantazaki, A. A. (2021). *In vitro* and *in silico* evaluation of the inhibitory effect of a curcumin-based oxovanadium (IV) complex on alkaline phosphatase activity and bacterial biofilm formation. *Applied Microbiology and Biotechnology*, 105(1), 147–168. <https://doi.org/10.1007/s00253-020-11004-0>
- Mombeshora, M., Chi, G. F., & Mukanganyama, S. (2021). Antibiofilm activity of extract and a compound isolated from *Triumfetta welwitschii* against *Pseudomonas aeruginosa*. *Biochemistry Research International*, 2021(1). <https://doi.org/10.1155/2021/9946183>
- Osland, A. M., Oastler, C., Konrat, K., Nesse, L. L., Brook, E., Richter, A. M., Gosling, R. J., Arvand, M., & Vestby, L. K. (2023). Evaluation of disinfectant efficacy against biofilm-residing wild-type *Salmonella* from the porcine industry. *Antibiotics*, 12(7), 1189. <https://doi.org/10.3390/antibiotics12071189>
- Paula, A. J., Hwang, G., & Koo, H. (2020). Dynamics of bacterial population growth in biofilms resemble spatial and structural aspects of urbanization. *Nature Communications*, 11(1), 1354. <https://doi.org/10.1038/s41467-020-15165-4>
- Raissa, G., Waturangi, D. E., & Wahjuningrum, D. (2020). Screening of antibiofilm and anti-quorum sensing activity of *Actinomyces* isolates extracts against aquaculture pathogenic bacteria. *BMC Microbiology*, 20(1), 343. <https://doi.org/10.1186/s12866-020-02022-z>
- Rubi, H., Mudey, G., & Kunjalwar, R. (2022). Catheter-associated urinary tract infection (CAUTI). *Cureus*, 14(10), e30385. <https://doi.org/10.7759/cureus.30385>
- Rueden, C. T., Schindelin, J., Hiner, M. C., DeZonia, B. E., Walter, A. E., Arena, E. T., & Eliceiri, K. W. (2017). ImageJ2: ImageJ for the next generation of scientific image data. *BMC Bioinformatics*, 18(1), 529. <https://doi.org/10.1186/s12859-017-1934-z>
- Ruhal, R., & Kataria, R. (2021). Biofilm patterns in gram-positive and gram-negative bacteria. *Microbiological Research*, 251, 126829. <https://doi.org/10.1016/j.micres.2021.126829>
- Schilcher, K., & Horswill, A. R. (2020). Staphylococcal biofilm development: structure, regulation,

- and treatment strategies. *Microbiology and Molecular Biology Reviews*, 84(3), e00026-19. <https://doi.org/10.1128/MMBR.00026-19>
- Schulze, K., López, D. A., Tillich, U. M., & Frohme, M. (2011). A simple viability analysis for unicellular cyanobacteria using a new autofluorescence assay, automated microscopy, and ImageJ. *BMC Biotechnology*, 11(1), 118. <https://doi.org/10.1186/1472-6750-11-118>
- JASP, T. (2024). *JASP*. <https://jasp-stats.org/>
- Vandeplassche, E., Coenye, T., & Crabbé, A. (2017). Developing selective media for quantification of multispecies biofilms following antibiotic treatment. *PLOS ONE*, 12(11), e0187540. <https://doi.org/10.1371/journal.pone.0187540>
- Yan, J., & Bassler, B. L. (2019). Surviving as a community: antibiotic tolerance and persistence in bacterial biofilms. *Cell Host & Microbe*, 26(1), 15–21. <https://doi.org/10.1016/j.chom.2019.06.002>
- Young, K., & Morrison, H. (2018). Quantifying microglia morphology from photomicrographs of immunohistochemistry prepared tissue using ImageJ. *Journal of Visualized Experiments*, 136, e57648. <https://doi.org/10.3791/57648-v>